



Heart Attack - Acute Myocardial Infarction (AMI)

Percent of Heart Attack Patients Given Aspirin at Arrival



Percent of Heart Attack Patients Given Aspirin at Discharge




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Heart Attack Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)



Percent of Heart Attack Patients Given Smoking Cessation Advice or Counseling




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Heart Attack Patients Given Beta Blocker at Discharge



Percent of Heart Attack Patients Given Fibrinolytic Medication Within 30 Minutes of Arrival




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Heart Attack Patients Given PCI (Percutaneous Coronary Interventions) Within 90 Minutes of Arrival



The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Quality Report Cards

Key Indicators	Saint Mary's Health Care	State Average	National Average	Top 10% of Hospitals Nationwide
Percent of Heart Attack Patients Given Aspirin at Arrival	100%	99%	99%	100%
Percent of Heart Attack Patients Given Aspirin at Discharge	96%	99%	99%	100%
Percent of Heart Attack Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	100%	96%	96%	100%
Percent of Heart Attack Patients Given Smoking Cessation Advice or Counseling	100%	100%	100%	100%
Percent of Heart Attack Patients Given Beta Blocker at Discharge	100%	99%	98%	100%
Percent of Heart Attack Patients Given Fibrinolytic Medication Within 30 Minutes of Arrival	N/A	59%	58%	100%
Percent of Heart Attack Patients Given PCI (Percutaneous Coronary Interventions) Within 90 Minutes of Arrival	100%	90%	91%	100%


The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.


State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Heart Failure

Percent of Heart Failure Patients Given Discharge Instructions



Percent of Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)



Percent of Heart Failure Patients Given Smoking Cessation Advice or Counseling



The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Quality Report Cards

Key Indicators	Saint Mary's Health Care	State Average	National Average	Top 10% of Hospitals Nationwide
Percent of Heart Failure Patients Given Discharge Instructions	95%	91%	90%	100%
Percent of Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function	99%	99%	98%	100%
Percent of Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD)	100%	95%	95%	100%
Percent of Heart Failure Patients Given Smoking Cessation Advice or Counseling	98%	99%	99%	100%


The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.


State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Pneumonia

Percent of Pneumonia Patients Assessed and Given Pneumococcal Vaccination



Percent of Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior to the Administration of the First Hospital Dose of Antibiotics




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Pneumonia Patients Given Smoking Cessation Advice or Counseling



Percent of Pneumonia Patients Given Initial Antibiotic(s) Within Six Hours After Arrival




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)



Percent of Pneumonia Patients Assessed and Given Influenza Vaccination



The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Quality Report Cards

Key Indicators	Saint Mary's Health Care	State Average	National Average	Top 10% of Hospitals Nationwide
Percent of Pneumonia Patients Assessed and Given Pneumococcal Vaccination	93%	93%	94%	100%
Percent of Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior to the Administration of the First Hospital Dose of Antibiotics	99%	96%	96%	100%
Percent of Pneumonia Patients Given Smoking Cessation Advice or Counseling	96%	98%	98%	100%
Percent of Pneumonia Patients Given Initial Antibiotic(s) Within Six Hours After Arrival	97%	96%	96%	100%
Percent of Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)	92%	94%	93%	100%
Percent of Pneumonia Patients Assessed and Given Influenza Vaccination	92%	91%	91%	100%


The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.


State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Surgical Care Improvement & Surgical Infection Prevention

Percent of Surgery Patients Who Received Preventative Antibiotic(s) One Hour Before Incision



Percent of Surgery Patients Who Received the Appropriate Preventative Antibiotic(s) for Their Surgery




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Surgery Patients Whose Preventative Antibiotic(s) Are Stopped Within 24 Hours After Surgery



Percent of All Heart Surgery Patients Whose Blood Sugar (Glucose) is Kept Under Good Control in the Days Right After Surgery




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Surgery Patients Needing Hair Removed From the Surgical Area Before Surgery, Who Had Hair Removed Using a Safer Method (Electric Clippers or Hair Removal Cream, Not a Razor)



Percent of Surgery Patients Whose Doctors Ordered Treatments to Prevent Blood Clots (Venous Thromboembolism) for Certain Types of Surgeries




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Percent of Surgery Patients Who Received Treatment to Prevent Blood Clots Within 24 Hours Before or After Selected Surgeries



Percent of Surgery Patients Who Were Taking Heart Drugs Called Beta Blockers Before Coming to the Hospital, Who Were Kept on the Beta Blockers During the Period Just Before and After Their Surgery




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.


Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Surgery Patients Whose Urinary Catheters Were Removed on the First or Second Day After Surgery



Outpatients Having Surgery Who Got an Antibiotic at the Right Time - Within One Hour Before Surgery




The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Outpatients Having Surgery Who Got the Right Kind of Antibiotic



The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.

Quality Report Cards

Key Indicators	Saint Mary's Health Care	State Average	National Average	Top 10% of Hospitals Nationwide
Percent of Surgery Patients Who Received Preventative Antibiotic(s) One Hour Before Incision	93%	97%	97%	100%
Percent of Surgery Patients Who Received the Appropriate Preventative Antibiotic(s) for Their Surgery	98%	98%	98%	100%
Percent of Surgery Patients Whose Preventative Antibiotic(s) Are Stopped Within 24 Hours After Surgery	97%	96%	96%	100%
Percent of All Heart Surgery Patients Whose Blood Sugar (Glucose) is Kept Under Good Control in the Days Right After Surgery	N/A	94%	94%	99%
Percent of Surgery Patients Needing Hair Removed From the Surgical Area Before Surgery, Who Had Hair Removed Using a Safer Method (Electric Clippers or Hair Removal Cream, Not a Razor)	99%	99%	100%	100%
Percent of Surgery Patients Whose Doctors Ordered Treatments to Prevent Blood Clots (Venous Thromboembolism) for Certain Types of Surgeries	100%	96%	95%	100%
Percent of Surgery Patients Who Received Treatment to Prevent Blood Clots Within 24 Hours Before or After Selected Surgeries	99%	95%	93%	100%
Percent of Surgery Patients Who Were Taking Heart Drugs Called Beta Blockers Before Coming to the Hospital, Who Were Kept on the Beta Blockers During the Period Just Before and After Their Surgery	92%	94%	94%	100%
Surgery Patients Whose Urinary Catheters Were Removed on the First or Second Day After Surgery	95%	91%	91%	100%
Outpatients Having Surgery Who Got an Antibiotic at the Right Time - Within One Hour Before Surgery	89%	90%	94%	100%
Outpatients Having Surgery Who Got the Right Kind of Antibiotic	91%	93%	95%	100%

The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

Saint Mary's Health Care data time period is January 2011 to July 2011.

State and National Averages and Top 10% of Hospitals Nationwide were obtained from the Hospital Compare website on October 25, 2011. The latest reporting period is from January 2010 through December 2010 for all indicators.